

## VU Research Portal

### High methane flux from an arctic floodplain (Indigirka lowlands, Eastern Siberia)

van Huissteden, J.; Maximov, T.C.; Dolman, A.J.

***published in***

Journal of Geophysical Research  
2005

***DOI (link to publisher)***

[10.1029/2005JG000010](https://doi.org/10.1029/2005JG000010)

***document version***

Publisher's PDF, also known as Version of record

[Link to publication in VU Research Portal](#)

***citation for published version (APA)***

van Huissteden, J., Maximov, T. C., & Dolman, A. J. (2005). High methane flux from an arctic floodplain (Indigirka lowlands, Eastern Siberia). *Journal of Geophysical Research*, 110(G02002).  
<https://doi.org/10.1029/2005JG000010>

**General rights**

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal ?

**Take down policy**

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

**E-mail address:**

[vuresearchportal.ub@vu.nl](mailto:vuresearchportal.ub@vu.nl)



## **High methane flux from an arctic floodplain (Indigirka lowlands, Eastern Siberia).**

J. van Huissteden, T.C. Maximov, A.J. Dolman

(1,3) Vrije Universiteit, Faculty of earth and Life Sciences, Department of Geo-environmental Sciences, De Boelelaan 1085, 1081 HV Amsterdam, The Netherlands, e-mail ko.van.huissteden@geo.falw.vu.nl

(2) Russian Academy of Sciences, Siberian Division, Institute for Biological Problems of Cryolithozone, 41, Lenin ave, Yakutsk, Sakha Republic, 677980 Russia, e-mail t.c.maximov@ibpc.ysn.ru

We measured methane fluxes from tundra soils on a river terrace and floodplain of a tributary of the Indigyrka river in Northeastern Siberia. The river terrace is underlain by shallow permafrost and shows a pattern of ice wedge polygons and a varied vegetation. The adjacent floodplain is characterized by frequent flooding, active sedimentation of silt and organics, a thicker vegetation and a more uniform vegetation of sedges and grasses. The methane fluxes measured on the river terrace compare well values found in other studies on tundra methane fluxes, with an average of  $4.3 \text{ mg.m}^{-2}.\text{hr}^{-1}$ , for wet sites  $7.2 \text{ mg.m}^{-2}.\text{hr}^{-1}$  and for dry sites  $0.18$ . Many dry sites show negative fluxes, down to  $-1.0 \text{ mg.m}^{-2}.\text{hr}^{-1}$ . The fluxes from floodplain sites are considerably higher, on average  $12.5 \text{ mg.m}^{-2}.\text{hr}^{-1}$ , for dry sites  $1.6 \text{ mg.m}^{-2}.\text{hr}^{-1}$  and for wet sites  $23.4 \text{ mg.m}^{-2}.\text{hr}^{-1}$ . We hypothesize that the large flux differences between river terrace and floodplain sites are caused by higher primary production of the floodplain vegetation, stimulated by nutrient and organic matter addition during flooding. Our results indicate that changes in fluvial discharge and organic matter transport of arctic rivers may have a considerable effect on arctic methane fluxes.